



UWS Academic Portal

Developmental social experience of parents affects behaviour of offspring in zebrafish

Tamilselvan, Priyadarshini; Sloman, Katherine

Published in:
Animal Behaviour

DOI:
[10.1016/j.anbehav.2017.09.009](https://doi.org/10.1016/j.anbehav.2017.09.009)

Published: 01/11/2017

Document Version
Peer reviewed version

[Link to publication on the UWS Academic Portal](#)

Citation for published version (APA):

Tamilselvan, P., & Sloman, K. (2017). Developmental social experience of parents affects behaviour of offspring in zebrafish. *Animal Behaviour*, 133, 153-160. <https://doi.org/10.1016/j.anbehav.2017.09.009>

General rights

Copyright and moral rights for the publications made accessible in the UWS Academic Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact pure@uws.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Developmental social experience of parents affects behaviour of offspring in zebrafish

P. Tamilselvan * and K.A. Sloman

School of Science and Sport, University of the West of Scotland, Paisley, U.K.

Received 7 March 2017

Initial acceptance 10 May 2017

Final acceptance 11 August 2017

MS number 17-00218

*Correspondence: P. Tamilselvan, School of Science and Sport, University of the West of Scotland, Paisley PA1 2BE, U.K.

E-mail address: priya.tamilselvan@uws.ac.uk

Interactions between conspecifics early in life have the potential to shape phenotypic differences between individuals. These changes in phenotype may subsequently be passed to future offspring, something that has been studied in live-bearing mammals where there is often an element of parental care. The present study considers the transgenerational effects of social environment in zebrafish, *Danio rerio*, an egg-laying animal that shows no parental care, thus removing any influence of parental interaction and allowing the effects of conspecific interaction to be clearly determined. Zebrafish (F₀) were reared from fertilization to reproduction in three different social treatments: isolation, groups of 30 or groups of 100. At 28 days post fertilization, individuals were tested for anxiety and activity and at 3 months old for aggression. These F₀ fish were raised to sexual maturity and bred within their treatment group. The F₁ generation were then raised in groups of 30, irrespective of parental social environment and were assessed for behaviour in the same way as their parents. Social isolation increased anxiety and decreased aggression in the F₀ fish compared to those raised in social groups of 100. F₀ fish raised in social groups of 30 showed an intermediate response. Differences in anxiety were not passed to the F₁ generation; however, offspring of socially

isolated fish were less aggressive than offspring of parents from social groups of 30 and 100.

The social environment that an individual experienced from fertilization to reproduction affected their own behaviour and the behaviour of their offspring.

Key words: activity, aggression, anxiety, behaviour, isolation, traits, transgenerational effects.

The early social environment experienced by an individual influences the development of behaviour as individuals in a social environment can learn from their conspecifics by observing them engaging in particular activities (Suboski & Templeton, 1989; Brown & Laland, 2001). Deprivation of social interaction early in development can, therefore, affect a variety of behaviours in a range of animals. For example, dairy calves housed individually immediately after birth were more reactive to environmental and social novelty than group-housed calves and calves housed with an older companion (Vieira, de Passille, & Weary, 2012). The majority of mammalian studies have considered the effects of social interactions after birth, although there is also evidence in rodents that interfetal communication can have a significant effect on behaviour later in life (vom Saal, 1989). In oviparous fish, which lay eggs into the external environment, exogenous cues such as the smell of predators or alarm cues from adults can alter developmental processes (Mirza, Chivers & Godin, 2001; Mourabit, Rundle, Spicer, & Sloman, 2010). For example, in rainbow trout, *Oncorhynchus mykiss*, raised in different social group sizes from fertilization, presence of conspecifics affected both physiology and behaviour. Trout raised in isolation had lower oxygen consumption rates and were less aggressive towards their own mirror image than individuals raised in social groups (Sloman & Baron, 2010).

51

52 The developmental environment experienced by an organism may alter not only its own
53 phenotype but also the behaviour of its offspring. Parental influence on offspring phenotype
54 (both maternal and paternal) can occur by genetic and epigenetic mechanisms. Maternal
55 effects are traditionally considered to be nongenomic, that is, they are not related to gene
56 sequence, although there is variation in the way 'maternal effects' are defined (Wolf & Wade,
57 2009). Both paternal and maternal effects have been documented in the literature, although
58 maternal effects have received the most attention and have been studied in a wide variety of
59 animals (mammals: Inhasz Kiss, Woodside, Felicio, Anselmo-Franci, & Damasceno, 2012;
60 birds: Guibert et al., 2011, Rubolini et al., 2005; reptiles: Robert, Vleck, & Bronikowski,
61 2009; Uller & Olsson, 2006; fish: Andersson, Silva, Steffensen, & Höglund, 2013; Eriksen et
62 al., 2011; Sloman, 2010). For example, a study on the pea aphid, *Acyrtosiphon pisum*,
63 showed that exposure of females to the alarm pheromone (E)- β -farnesene prior to
64 reproduction, a cue for predation risk, resulted in a change of feeding sites in offspring
65 (Keiser & Mondor, 2013). Postnatal communal rearing in Balb/c mice, *Mus musculus*,
66 induced transgenerational effects on emotional and reproductive behaviour of offspring
67 (Curley, Davidson, Bateson, & Champagne, 2009). Communally raised females that received
68 more postpartum maternal care exhibited lower anxiety, built higher quality nests and showed
69 more postpartum care as adults than standard reared females. This behaviour was further
70 carried through to F₂ mice that exhibited lower anxiety, larger litter size and increased
71 nursing, suggesting the effect of postnatal social environment on the behaviour of offspring
72 across generations (Curley et al., 2009).

73

74 Early in development the epigenome of an individual can be influenced by environmental and
75 nutritional factors, such as transfer of hormones or nutrient provision by mothers to eggs or

offspring (Nafee, Farrell, Carroll, Fryer, & Ismail, 2007). Parental conditions can alter the phenotype of offspring (Chen et al., 2013; Franzke & Reinhold, 2013; Krause & Naguib, 2014; Pittet, Le Bot, Houdelier, Richard-Yris, & Lumineau, 2014) and while some authors may conclude that these transgenerational effects occur via epigenetic mechanisms (Youngson & Whitelaw, 2008), others would argue that when considering environmentally-induced effects, an epigenetic basis can be inferred only if changes last over multiple generations (Grossniklaus, William, Ferguson-Smith, Pembrey, & Lindquist, 2013; see also Burggren, 2014). Mechanisms that allow transfer of information about the maternal environment to offspring are likely to be advantageous if they prepare the offspring for the environment they will be born into (Love, McGowan, & Sheriff, 2013; Sheriff & Love, 2013). Parental mechanisms that allow adjustments of offspring phenotype based on immediate parental environment may allow more flexibility than selection on genotypes (Crews, 2008; Wisenden, Sailer, Radenic, & Sutrisno, 2011).

A number of studies have looked at the effect of variations in the social environment during development. However, it is not known whether changes in phenotype induced by early social interactions can be passed across generations. Furthermore, previous studies have addressed transgenerational effects wherein the parents were exposed to the experimental manipulation only during a certain point of their own development and transferred to control conditions before reaching sexual maturity; any responses in offspring could thus be the result of differences in natal and adult environments (reviewed in Burton & Metcalfe, 2014). Therefore, the aim of the present study was, first, to examine the effects of different social environments maintained from fertilization to reproduction in F_0 zebrafish, *Danio rerio*, and, second, to investigate any transfer of behavioural effects to the subsequent F_1 generation. Social environments were varied by number of individuals rather than stocking density; it

was hypothesized that zebrafish raised in different social environments would exhibit differences in behaviour later in life and these differences in behavioural phenotype may be transmitted to future generations.

Methods

Adult zebrafish (AB, TL mixed strains) from an existing stock at the University of the West of Scotland were held on a recirculating system (27 ± 1 °C; pH 7.1 ± 0.4 ; dissolved oxygen $90\pm5\%$; 14:10 h light:dark) and fed *Artemia* or Aquarian tropical flake twice daily. Fish were bred to produce embryos (F_0 generation) which were collected within 30 min of fertilization and placed into group sizes of 1 ($N=36$), 30 ($N=3$) and 100 ($N=3$). The different group sizes were held in 50, 500 and 5000 ml containers with 10, 30 and 1000 ml of system water, respectively, and placed in a water bath at 28.5 °C. The walls of the containers were opaque to block any visual cues and a 30% water change was carried out daily. From 5 days post fertilization (dpf) larvae were fed Liquifry and ZM 00 daily. One week after hatching, fry were transferred in their group sizes into flow-through tanks with opaque sides on the main recirculating system. Tanks sizes were 1, 3 and 12 litres containing 0.94, 2.95 and 11.5 litres of water, respectively. Fry were fed ZM 100 from 11 dpf and ZM 200 and *Artemia* from 15 dpf. From 30 dpf they were fed twice daily with flake food and *Artemia*. Variations in tank sizes between the different treatments ensured that treatments represented a change in number of individuals within the social group, not differences in stocking density. An overview of the experimental design is shown in Fig. 1.

Breeding of F_0 fish

When the F_0 fish were 3 months old, they were bred to produce the F_1 generation. To achieve this, F_0 fish raised in social isolation were combined within their treatment resulting in three replicate breeding tanks for each treatment. Thirty F_1 embryos from each replicate of each F_0 treatment (i.e. $N=3$) were then raised in 1.5-litre containers held at 28.5 °C. Thus, F_1 offspring from all F_0 treatments were held at the same density and the only difference between F_1 treatments was the social environment experienced by their parents. Two weeks after hatching, fry were transferred into 3-litre flow-through tanks on the main recirculating system. The feeding regime was the same as for the F_0 generation.

Behavioural testing of F_0 and F_1 fry

At 28 dpf F_0 ($N=27$) and F_1 ($N=18$) fry were tested for their behaviour in the light and dark box test and the novel tank diving test. At 3 months old, fish ($N=27$ F_0 ; $N=18$ F_1) were tested for their response to their own mirror image as zebrafish show similar levels of aggression towards a mirror image as towards an opponent (Ariyomo & Watt, 2013).

Light and Dark Box Test

The experimental tank (16 x 10 cm and 10.5 cm high) was divided vertically in half and covered externally on the sides and bottom by white or black paper. The top was left completely open. The half of the tank covered externally with a white background represented the light compartment and the half with a black background was the dark compartment. An opaque cylinder was placed in the centre of the tank creating a compartment (5 cm diameter) for acclimating the fish. The tank was filled with system water which was replaced at the end of every trial (Blaser & Rosemberg, 2012; Maximino et al., 2010a; Maximino, de Brito, Dias, Gouveia, & Morato, 2010b). Fish were individually placed

into the central compartment for 5 min, after which time the walls of the central compartment were removed, allowing the fish to freely explore the tank. Behaviour was recorded for 10 min with a video camera (Panasonic SD video camera, SDR-S50) placed above the arena. The tank was illuminated with a white daylight bulb (70 W) and the illumination kept constant between trials. The behavioural end points measured were the proportion of time the fish spent in the dark compartment (scototaxis), the time spent beside the walls (thigmotaxis) and the number of transitions between the light and dark compartment (activity) (Blaser & Rosemberg, 2012; Maximino et al., 2010a, b).

Novel Tank Diving Test

A trapezoidal tank, greater in horizontal cross-sectional area at the top than the bottom, (top: 16 x 10 cm; bottom: 15 x 9 cm; 10.5 cm deep) was visually divided horizontally on the outside, half way between the top water line (8 cm) and the bottom of the tank (Cachat et al., 2010; Egan et al., 2009; Levin, Bencan, & Cerutti, 2007). The tank was filled with system water which was replaced at the end of every trial. Individual fish were transferred to the experimental tank and behaviour was recorded for 6 min with a video camera. The behavioural parameters measured were latency to enter the upper compartment of the tank, time spent in the upper and lower compartments of the tank and number of entries into the upper and lower compartments (activity). A longer latency to reach the upper compartment and surface and reduced entries into the upper compartment are considered to be indicators of stress and anxiety (Egan et al., 2009; Levin et al., 2007).

Mirror Image Test

Individual fish were placed in a test tank (16 x 10 cm and 10.5 cm high) containing a covered mirror and left to acclimate for 20 h. Before commencement of behavioural recordings, the mirror was revealed for 1 min to allow the fish to get used to the disturbance caused by revealing the mirror. After a further 15 min, the mirror was revealed once again and behaviour was video recorded for 1 h. The behavioural parameters measured were latency to first attack the mirror, frequency of mirror biting and time spent within 5 cm of the mirror.

Ethical Note

All experiments adhered to the ASAB/ABS guidelines for the use of animals in research and the U.K. Home Office guidelines (licence: 70/8539). All fish were held in appropriate water quality (as documented above) and water quality was checked daily for all treatments. All experiments were mild and did not cause significant pain or lasting harm to the fish; a mirror image test was used to measure aggression rather than pairing of conspecifics which can lead to physical damage.

Data analysis

All the behavioural videos were observed blind, that is, the observer did not know which treatment groups they came from. Data analysis was carried out using the statistical software SPSS Statistics 20 (IBN, Armon, NY, U.S.A.). Data were tested for normality using the Kolmogorov–Smirnov test and found to be normally distributed. Analysis of variance was used to compare behaviour between the three treatment groups in the F₀ and F₁ generations, respectively. Post hoc testing with a least significant difference test was performed on the behavioural parameters that showed significant results ($P < 0.05$) and is presented in the figures. For the F₁ generation, tank replicate was included as a random effect in the model.

Results

Light and Dark Box Test

For the F_0 generation, fish raised in social groups of 100 displayed significantly less scototaxis ($F_{2, 66} = 4.891$, $P=0.010$; Fig. 2a), spending less time in the dark compartment, than those reared in groups of 30 or in social isolation. Fish raised in isolation displayed significantly more activity ($F_{2, 66} = 4.783$, $P=0.011$; Fig. 2b) than those in social groups of 100. Time spent beside the walls (thigmotaxis) was not significantly different between treatments ($F_{2, 66} = 0.310$, $P=0.735$). Neither was there any significant difference in thigmotaxis between treatments when data were analysed separately for the light and dark compartments (light: $F_{2, 66} = 0.602$, $P=0.179$; dark: $F_{2, 66} = 2.129$, $P=0.076$). In the F_1 generation, there were no significant effects of treatment on scototaxis ($F_{2, 51} = 0.183$, $P=0.840$), activity ($F_{2, 51} = 1.528$, $P=0.321$) or thigmotaxis ($F_{2, 51} = 0.399$, $P=0.695$).

Novel Tank Diving Test

In the F_0 generation, there was a reduced latency to reach the upper compartment ($F_{2, 64} = 3.563$, $P=0.034$; Fig. 3a) in the fish raised in groups of 100 compared to those from isolation. However, time spent in the upper or lower compartment was not significantly different ($F_{2, 64} = 0.949$, $P=0.392$) between treatments. Fish raised in isolation displayed significantly more activity ($F_{2, 64} = 10.597$, $P<0.001$; Fig. 3b) than those raised in social groups. For the F_1 generation, there was no significant effect of treatment on latency to reach the upper compartment ($F_{2, 51} = 0.674$, $P=0.559$), time spent in the upper compartment ($F_{2, 51} = 0.175$, $P=0.845$) or activity ($F_{2, 51} = 1.990$, $P=0.251$).

Mirror Image Test

In the F_0 generation, fish raised in social isolation displayed significantly fewer attacks towards the mirror than those raised in social groups of 100 and 30 ($F_{2, 73} = 3.626$, $P=0.032$; Fig. 4a). There was no difference in latency to attack the mirror ($F_{2, 73} = 1.699$, $P=0.190$) or the time spent beside the mirror ($F_{2, 73} = 1.986$, $P=0.145$). In the F_1 generation the latency of fish to attack their own mirror image was not significantly different between F_1 fish whose parents were raised in different social environments ($F_{2, 51} = 0.425$, $P=0.680$). However, F_1 fish from parents raised in social isolation displayed significantly fewer attacks ($F_{2, 51} = 142.039$, $P=0.007$; Fig. 4b) and spent less time beside the mirror ($F_{2, 51} = 322.775$, $P<0.001$; Fig. 4c) than those from parents raised in social groups.

Discussion

The present study has shown that the social environment experienced by zebrafish from fertilization to reproduction affects anxiety and aggression later in life and that differences in aggression are passed on to offspring regardless of their rearing environment. To our knowledge, this is the first study in fish to have considered whether behavioural changes in the F_0 generation induced by their social environment can be passed on to their offspring. In fish, social isolation has been shown to alter behaviour (Ichihashi, Ichikawa, & Matsushima, 2004; Gomez-Laplaza & Morgan, 2000; Zellner, Padnos, Hunter, MacPhail, & Padilla, 2011); however, most of the previous studies in fish have manipulated the social environment after hatching, whereas in the present study zebrafish embryos were separated immediately after fertilization.

In the present study, in the F₀ generation, individuals raised in either social isolation or in groups of 30 were more scototactic (spent more time in the dark) than those raised in groups of 100. Scototaxis as a measure of anxiety in zebrafish has been validated in a number of previous studies (Blaser & Rosemberg, 2012; Maximino et al., 2010a,b), anxious fish being more scototactic (Maximino et al., 2012; Maximino et al., 2010a). It has been suggested that scototaxis is a form of defence where camouflaging with the substratum can help avoid predation (Maximino et al., 2010b). Thigmotaxis (wall hugging) is another indicator of anxiety. This phenomenon has been found in mice and it is proposed to assist in search for shelter and protection (Treit & Fundytus, 1988). Anxious fish exhibit higher thigmotaxis (Blaser & Gerlai, 2006; Champagne, Hoefnagels, de Kloet & Richardson, 2010; Maximino et al., 2010a) but in the present study F₀ fish showed no difference in thigmotaxis. As previous studies have shown that more thigmotaxis can occur in the dark (Blaser, Chadwick, & McGinnis, 2010), the thigmotaxis data were further analysed for the dark and light compartments separately but no significant effects were found. Therefore, one anxiety measure (scototaxis) indicated that the fish raised in isolation and small social groups were more anxious, whereas the second measure of anxiety (thigmotaxis) found no effects. These results are in line with studies by Blaser et al. (2010) who also found thigmotaxis behaviours difficult to interpret. They suggested that 'thigmotaxis' includes a greater variety of behaviours than simply 'proximity to the walls' and therefore may require additional behavioural measurements to relate to anxiety. The effects on anxiety levels indicated by differences in scototaxis in the light and dark box test, however, were further substantiated by the results of the novel tank diving test where anxious fish take longer to reach the upper part of the tank (Egan et al., 2009; Gerlai, 2003; Levin et al., 2007). Fish raised in isolation were more anxious than those raised in groups of 100 as they showed a longer latency to reach the upper compartment, with fish raised in groups of 30 showing an intermediate response.

270 In both the light and dark box test and the novel tank diving test, fish raised in social isolation
271 were more active than those raised in social groups of 100. In the light and dark box test, fish
272 raised in social groups of 30 were more active than those raised in groups of 100, whereas in
273 the novel tank diving test fish raised in groups of 30 showed a more intermediate response.
274 Test-specific differences in the behaviour of zebrafish have been reported in previous studies
275 (Blaser & Gerlai, 2006); however, the overall pattern in both tests shows an increased level of
276 activity with decreasing social experience. Activity has previously been suggested as a
277 measure of anxiety (Blaser et al., 2010; Maximino et al., 2010a; Peng et al., 2016); therefore,
278 the increased activity seen in the fish raised in social isolation supports the higher anxiety
279 suggested by scototaxis and latency to reach the upper compartment.

280

281 None of the behavioural parameters measured in either the light and dark box test or the
282 novel tank diving test were significantly different in the F_1 generation where offspring of F_0
283 fish reared in social isolation and social groups of 30 and 100 were all reared in groups of 30.
284 This would suggest that although the F_0 socially isolated fish were more anxious, these
285 effects were not passed on to their offspring. The fact that differences in anxiety were not
286 transferred to offspring may be because anxiety can be a disadvantageous trait to pass on as it
287 can lead to increased predation (McGhee & Bell, 2014). In the present study, the genetic pool
288 was assumed to be equally distributed between treatments at the start, as we were specifically
289 interested in nongenomic effects of parental social environment. As both mothers and fathers
290 were exposed to the same social environments, we cannot separate maternal from paternal
291 effects. There is evidence that some behavioural traits such as boldness and aggression have a
292 heritable component (Brown, Burgess, & Braithwaite, 2007; Ariyomo, Carter, & Watt, 2013),

and further studies on the heritability of anxiety, and potential interactions with parental effects, are warranted.

At approximately 100 dpf, F₀ zebrafish raised in different social environments displayed differences in their response to their own mirror image. The mirror image test was used to measure aggression as it represents a measure of how individuals would interact with conspecifics (Moretz, Martins, & Robison, 2007). Fish are unable to recognize their own image and respond to the mirror image as if it were an intruder (Rowland, 1999). Aggression is exhibited by both sexes in zebrafish (Moretz et al., 2007) and is important in reproductive success and establishment of dominance hierarchies (Paull, Filby, Giddins, Coe, Hamilton, & Tyler., 2010). Although the mirror image test is generally considered to be good for measuring some behavioural parameters over others (Balzarini, Taborsky, Wanner, Kock, Felizia, & Frommen, 2014; Elwood, Stoilova, McDonnell, Earley, & Arnott, 2014), Ariyomo and Watt (2013) found that there were no differences in the rate of aggression of zebrafish towards a mirror image and an opponent, thus demonstrating the validity of the mirror image test as an indicator of aggression in zebrafish. Overall, fish raised in social groups of 30 and 100 were more aggressive than those held in social isolation. The results of the present study are similar to results obtained by Sloman and Baron (2010) who found that rainbow trout raised in social isolation performed half as many bites as those raised in social groups. In both their study and ours, fish were raised in their respective social conditions from fertilization until they were used in the mirror image test, so those raised in social isolation had never encountered another fish. In the present study, there was no difference between treatments in latency to attack the mirror or the time spent beside the mirror, suggesting an innate fighting ability of fish raised in social isolation.

It is possible that the absence of conspecifics during the very early stages of development might have led to adults with reduced aggressive behaviour since they were not receiving cues about the presence of conspecific competition. Sloman and Baron (2010) suggested that the higher prehatching metabolic rate of embryos raised in social groups indicated their ability to detect cues from conspecifics in their environment during development and through their higher metabolic rate have an increased competitive ability at first feeding. Similarly, it is likely that in the present study embryos raised in social groups could detect cues from conspecifics during development which increased aggression, reinforced by social interaction after hatching. In social groups individuals learn from their conspecifics (Suboski & Templeton, 1989) whereas fish raised in social isolation had no opportunity to learn. The present study confirms that fish raised in social groups are more aggressive and here we found that this behaviour is passed to their offspring. This is likely to be advantageous for individuals born into a competitive environment. In contrast, social isolation appears to reduce aggression, which persists in the next generation even when offspring are raised in social groups. In this scenario, where there is a mismatch between parent and offspring environment, reduced aggression levels may be disadvantageous. To examine this further, it would be interesting to know how long social isolation needs to persist in the F_0 generation for it to be transferred to the next generation. For example, if an embryo is separated from its conspecifics from fertilization until hatching only, does it recover normal levels of aggression? Also, it would be interesting to know whether the higher aggression seen in the F_1 generation whose parents were raised in social groups persists if the F_1 generation are all reared in social isolation. Aggression levels were much higher in the F_1 than in the F_0 generation (Fig. 4). At present, we have no clear explanation for this difference and future studies are required to see if this is maintained across further generations.

The advantages of a mother transferring information about her environment to her offspring, mostly associated with maternal transfer of stress hormones, have been reviewed previously (Love et al., 2013; Sheriff & Love, 2013). For example, female sticklebacks, *Gasterosteus aculeatus*, can transfer information about the level of predation in their environment to their offspring, and it has been suggested that this occurs via cortisol transfer (Giesing, Suski, Warner, & Bell, 2011). Eaton, Edmonds, Henry, Snellgrove, and Sloman (2015) speculated that the lack of associative learning by guppy offspring of mildly stressed mothers may be advantageous. If offspring are to be born into an unpredictable environment then a lack of associative learning, or ability to forget, may be adaptive (Warburton, 2003). The mechanisms by which differences in aggression levels are transferred between generations in the present study remain unknown but it is possible that different social environments resulted in different levels of cortisol or nutrients being deposited in eggs during oogenesis.

In conclusion, offspring of fish reared in social isolation were less aggressive than the offspring of group-reared fish. However, there were no significant differences in behavioural parameters related to anxiety and activity in the F₁ generation. These results suggest that zebrafish can transfer information about their environment to offspring, leading to nongenomic transgenerational transmission of behavioural phenotypes. The mechanisms underlying these differences are not clear and future studies should look both at whether these effects persist across multiple generations and for evidence of epigenetic mechanisms.

Acknowledgments

We thank the technicians Deirdre Galbraith and Edith Burns for their help with animal husbandry and two anonymous reviewers for comments on a previous draft.

367

368 **References**

369 Ariyomo, T. O., & Watt, P.J. (2013). Aggression and sex differences in lateralization in the

370 zebrafish. *Animal Behaviour*, 86, 617-622.

371 Ariyomo, T.O., Carter, M., & Watt, P.J. (2013). Heritability of boldness and aggressiveness

372 in the zebrafish. *Behavioural Genetics*, 43, 161-167.

373 Balzarini, V., Taborsky, M., Wanner, S., Koch, F., & Frommen, J.G. (2014). Mirror, mirror

374 on the wall: the predictive value of mirror tests for measuring aggression in fish.

375 *Behaviour Ecology and Sociobiology*, 68, 871-878

376 Blaser, R., & Gerlai, R. (2006). Behavioral phenotyping in zebrafish: Comparison of three

377 behavioral quantification methods. *Behavior Research Methods*, 38(3), 456-469.

378 doi:10.3758/bf03192800

379 Blaser, R. E., Chadwick, L., & McGinnis, G. C. (2010). Behavioral measures of anxiety in

380 zebrafish (*Danio rerio*). *Behavioural Brain Research*, 208(1), 56-62.

381 doi:10.1016/j.bbr.2009.11.009

382 Blaser, R. E., & Rosemberg, D. B. (2012). Measures of anxiety in zebrafish (*Danio rerio*):

383 Dissociation of black/white preference and novel tank test. *Plos One*, 7(5).

384 doi:10.1371/journal.pone.0036931

385 Brown, C., & Laland, K. (2001). Social learning and life skills training for hatchery reared

386 fish. *Journal of Fish Biology*, 59, 471-493. 10.1111/j.1095-8649.2001.tb02354.x

387 Brown, C., Burgess, F., & Braithwaite, V. A. (2007). Heritable and experiential effects on

388 boldness in a tropical poeciliid. *Behavioral Ecology and Sociobiology*, 62(2), 237-243.

389 doi:10.1007/s00265-007-0458-3

- Burggren, W.W. (2014). Epigenetics as a source of variation in comparative animal physiology – or – Lamarck is lookin’ pretty good these days. *Journal of Experimental Biology*, 217, 682-689. 10.1242/jeb.086132
- Burton, T., & Metcalfe, N. B. (2014). Can environmental conditions experienced in early life influence future generations? *Proceedings of the Royal Society B*, 281(1785), 20140311-20140311. doi:10.1098/rspb.2014.0311
- Andersson, M. A., Silva, P.I.M., Steffensen, J.F., & Höglund, E. (2013). Effects of maternal stress coping style on offspring characteristics in rainbow trout (*Oncorhynchus mykiss*). *Hormones and Behavior*, 63, 674-674.
- Cachat, J., Canavello, P., Elegante, M., Bartels, B., Hart, P., Bergner, C., . . . Kalueff, A.V. (2010). Modelling withdrawal syndrome in zebrafish. *Behavioural brain research*, 208(2), 371-376. Doi:10.1016/j.bbr.2009.12.004
- Champagne, D. L., Hoefnagels, C.C.M., de Kloet, R.E., & Richardson, M.K. (2010). Translating rodent behavioral repertoire to zebrafish (*Danio rerio*): Relevance for stress research. *Behavioural Brain Research*, 214, 332-342.
- Chen, J., Das, S.R., La Du, J., Corvi, M.M., Bai, C., Chen, Y.,& Huang, C. (2013). Chronic PFOS exposures induce life stage-specific behavioral deficits in adult zebrafish and produce malformation and behavioral deficits in F1 offspring. *Environmental Toxicology and Chemistry*, 32, 201-206.
- Crews, D. (2008). Epigenetics and its implications for behavioural neuroendocrinology. *Frontiers in Neuroendocrinology*, 29, 344-357.
- Curley, J. P., Davidson, S., Bateson, P., & Champagne, F. A. (2009). Social enrichment during postnatal development induces transgenerational effects on emotional and reproductive behavior in mice. *Frontiers in Behavioral Neuroscience*, 3. doi:10.3389/neuro.08.025.2009

415 Eaton, L., Edmonds, E.J., Henry, T.B., Snellgrove, D.L., & Sloman, K.A. (2015). Mild
 416 maternal stress disrupts associative learning and increases aggression in offspring.
 417 *Hormones and Behavior*, 71, 10-15.

418 Egan, R. J., Bergner, C. L., Hart, P. C., Cachat, J. M., Canavello, P. R., Elegante, M. F.,
 419 & Kalueff, A. V. (2009). Understanding behavioral and physiological
 420 phenotypes of stress and anxiety in zebrafish. *Behavioural Brain Research*, 205(1), 38-
 421 44. doi:10.1016/j.bbr.2009.06.022

422 Elwood, R.W., Stoilova, V., McDonnell, A., Earley, R.L., & Arnott, G. (2014). Do mirrors
 423 reflect reality in agonistic encounters? A test of mutual cooperation in displays. *Animal*
 424 *Behaviour*, 97, 63-67.

425 Eriksen, M. S., Faerevik, G., Kittilsen, S., McCormick, M.I., Damsgard, B., Braithwaite,
 426 V.A., & Bakken, M.A. (2011). Stressed mothers - troubled offspring: a study of
 427 behavioural maternal effects in farmed *Salmo salar*. *Journal of Fish Biology*, 79, 575-
 428 586.

429 Franzke, A., & Reinhold, K. (2013). Transgenerational effects of diet environment on life-
 430 history and acoustic signals of a grasshopper. *Behavioural Ecology*, 24, 734-739.

431 Gerlai, R. (2003). Zebrafish: An uncharted behavior genetic model. *Behavior Genetics*, 33(5),
 432 461-468. doi:10.1023/a:1025762314250

433 Giesing, E.R., Suski, C.D., Warner, R.E., & Bell, A.M. (2011). Female sticklebacks transfer
 434 information via eggs: effects of maternal experience with predators on offspring.
 435 *Proceedings of the Royal Society B*, 278, 1753-1759.

436 Gomez-Laplaza, L. M., & Morgan, E. (2000). Laboratory studies of the effects of short-term
 437 isolation on aggressive behaviour in fish. *Marine and Freshwater Behaviour and*
 438 *Physiology*, 33, 63-102.

439 Grossniklaus, U., William, K.G., Ferguson-Smith, A.C., Pembrey, M., & Lindquist, S.
 440 (2013). Transgenerational epigenetic inheritance: how important is it? *Nature Review*
 441 *Genetics*, 14, 228.

442 Guibert, F., Richard-Yris, M.-A., Lumineau, S., Kotrschal, K., Bertin, A., Petton, C., &
 443 Houdelier, C. (2011). Unpredictable mild stressors on laying females influence the
 444 composition of Japanese quail eggs and offspring's phenotype. *Applied Animal*
 445 *Behaviour Science*, 132, 51-60.

446 Ichihashi, T., Ichikawa, Y., & Matsushima, T. (2004). A non-social and isolate rearing
 447 condition induces an irreversible shift toward continued fights in the male fighting
 448 fish (*Betta splendens*). *Zoological Science*, 21, 723-729.

449 Inhasz Kiss, A. C., Woodside, B., Felicio, L.F., Anselmo-Franci, J. & Damasceno, D.C.
 450 (2012). Impact of maternal mild hyperglycemia on maternal care and offspring
 451 development and behavior of Wistar rats. *Physiology & Behavior*, 107, 292-300.
 452 [dx.doi.org/10.1016/j.physbeh.2012.08.001](https://doi.org/10.1016/j.physbeh.2012.08.001)

453 Keiser, C. N., & Mondor, E. B. (2013). Transgenerational behavioral plasticity in a
 454 parthenogenetic insect in response to increased predation risk. *Journal of Insect*
 455 *Behavior*, 26(4), 603-613. doi:10.1007/s10905-013-9376-6

456 Krause, E. T., & Naguib, M. (2014). Effects of parental and own early developmental
 457 conditions on the phenotype in zebra finches (*Taeniopygia guttata*). *Evolutionary*
 458 *Ecology*, 28, 263-275.

459 Levin, E. D., Bencan, Z., & Cerutti, D. T. (2007). Anxiolytic effects of nicotine in zebrafish.
 460 *Physiology & Behavior*, 90(1), 54-58. doi:10.1016/j.physbeh.2006.08.026

461 Love, O.P., McGowan, P., & Sheriff, M.J. (2013). Maternal adversity and ecological
 462 stressors in natural populations: the role of stress axis programming in individuals, with
 463 implications for populations and communities. *Functional Ecology*, 27, 81-92.

464 Maximino, C., Benzecry, R., Oliveira, K. R. M., Batista, E. D. O., Herculano, A. M.,
 465 Rosemberg, D. B., & Blaser, R. (2012). A comparison of the light/dark and
 466 novel tank tests in zebrafish. *Behaviour*, 149(10-12), 1099-1123.
 467 doi:10.1163/1568539x-00003029

468 Maximino, C., de Brito, T. M., Colmanetti, R., Pontes, A. A. A., de Castro, H. M., de
 469 Lacerda, R. I. T., & Gouveia, A. (2010a). Parametric analyses of anxiety in
 470 zebrafish scototaxis. *Behavioural Brain Research*, 210(1), 1-7.
 471 doi:10.1016/j.bbr.2010.01.031

472 Maximino, C., de Brito, T. M., Dias, C., Gouveia, A., & Morato, S. (2010b). Scototaxis as
 473 anxiety-like behavior in fish. *Nature Protocols*, 5(2), 209-216.
 474 doi:10.1038/nprot.2009.225

475 McGhee, K. E., & Bell, A. M. (2014). Paternal care in a fish: epigenetics and fitness
 476 enhancing effects on offspring anxiety. *Proceedings of the Royal Society B-Biological*
 477 *Sciences*, 281(1794). doi:10.1098/rspb.2014.1146

478 Mirza, R.S., Chivers, D.P., & Godin, J.G.J. (2001). Brook charr alevins alter timing of next
 479 emergence in response to chemical cues from fish predators. *Journal of Chemical*
 480 *Ecology*, 27, 1775-1785.

481 Moretz, J. A., Martins, E. P., & Robison, B. D. (2007). Behavioral syndromes and the
 482 evolution of correlated behavior in zebrafish. *Behavioral Ecology*, 18(3), 556-562.
 483 doi:10.1093/beheco/arm011

484 Mourabit, S., Rundle, S.D., Spicer, J.I., & Sloman, K.A. (2010). Alarm substance from adult
 485 zebrafish alters early embryonic development in offspring. *Biology Letters*, 6, 525-528.

486 Nafee, T.M., Farrell, W.E., Carroll, W.D., Fryer, A.A., & Ismail, K.M.K. (2008). Epigenetic
 487 control of fetal gene expression. *BJOG*, 115, 158-168. 10.1111/j.1471-
 488 0528.2007.01528.x

489 Paull, G. C., Filby, A. L., Giddins, H. G., Coe, T. S., Hamilton, P. B., & Tyler, C. R. (2010).
 490 Dominance hierarchies in zebrafish (*Danio rerio*) and their relationship with
 491 reproductive success. *Zebrafish*, 7(1), 109-117. doi:10.1089/zeb.2009.0618

492 Peng, X. L., Lin, J., Zhu, Y. D., Liu, X. Y., Zhang, Y. L., Ji, Y. X., Y..... & Li,
 493 Q. (2016). Anxiety-related behavioral responses of pentylenetetrazole-treated
 494 zebrafish larvae to light-dark transitions. *Pharmacology, Biochemistry and Behavior*,
 495 145, 55-65. doi:10.1016/j.pbb.2016.03.010

496 Pittet, F., Le Bot, O., Houdelier, C., Richard-Yris, M., & Lumineau, S. (2014). Motherless
 497 quail mothers display impaired maternal behavior and produce more fearful and less
 498 socially motivated offspring. *Developmental Psychobiology*, 56, 622-634.

499 Robert, K. A., Vleck, C., & Bronikowski, A.M. (2009). The effects of maternal
 500 corticosterone levels on offspring behavior in fast- and slow-growth garter snakes
 501 (*Thamnophis elegans*). *Hormones and Behavior*, 55, 24-32.

502 Rowland, W. J. (1999). Studying visual cues in fish behavior: a review of ethological
 503 techniques. *Environmental Biology of Fishes*, 56(3), 285-305.
 504 doi:10.1023/a:1007517720723

505 Rubolini, D., Romano, M., Boncoraglio, G., Ferrari, R.P., Martinelli, R., Galeotti, P.,
 506 & Saino, N. (2005). Effects of elevated egg corticosterone levels on behavior,
 507 growth, and immunity of yellow-legged gull (*Larus michahellis*) chicks. *Hormones and*
 508 *Behavior*, 47, 592-605.

509 Sheriff, M.J., & Love, O.P. (2013). Determining the adaptive potential of maternal stress.
 510 *Ecology Letters*, 16, 271-280.

511 Sloman, K. A. (2010). Exposure of ova to cortisol pre-fertilisation affects subsequent
 512 behaviour and physiology of brown trout. *Hormones and Behavior* 58, 433-439.

513 Sloman, K. A., & Baron, M. (2010). Conspecific presence affects the physiology and
 514 behaviour of developing trout. *Physiology & Behavior*, 99(5), 599-604.
 515 doi:10.1016/j.physbeh.2010.01.027

516 Suboski, M. D., & Templeton, J. J. (1989). Life skills training for hatchery fish - social -
 517 learning and survival. *Fisheries Research*, 7(4), 343-352. doi:10.1016/0165-
 518 7836(89)90066-0

519 Treit, D., & Fundytus, M. (1988). Thigmotaxis as a test for anxiolytic activity in
 520 rats. *Pharmacology Biochemistry and Behavior*, 31(4), 959-962. doi: 10.1016/0091-
 521 3057(88)90413-3

522 Uller, T., & Olsson, M. (2006). Direct exposure to corticosterone during embryonic
 523 development influences behaviour in an ovoviviparous lizard. *Ethology*, 112, 390-397.

524 Vieira, A. D., de Passille, A.M., & Weary, D.M. (2012). Effects of the early social
 525 environment on behavioral responses of dairy calves to novel events. *Journal of Dairy*
 526 *Science*, 95, 5149-5155. dx.doi.org/10.3168/jds.2011-5073

527 vom Saal, F.S. (1989). Sexual differentiation in litter-bearing mammals: influence of sex of
 528 adjacent fetuses in utero. *Journal of Animal Science*, 67, 1824-1840.

529 Warburton, K. (2003). Learning of foraging skills by fish. *Fish and Fisheries*, 4, 203-215.

530 Wisenden, B. D., Sailer, C.D., Radenic, S.J., & Sutrisno, R. (2011). Maternal inheritance and
 531 exploratory-boldness behavioural syndrome in zebrafish. *Behaviour*, 148, 1443-1456.

532 Wolf, J.B., & Wade, M.J. (2009). What are maternal effects (and what are they not)?
 533 *Philosophical Transactions of the Royal Society*, 364, 1107-1115.
 534 10.1098/rstb.2008.0238

535 Youngson, N.A., & Whitelaw, E. (2008). Transgenerational epigenetic effects. *Annual*
 536 *Review of Genomics and Human Genetics*, 9, 233-257.

537 Zellner, D., Padnos, B., Hunter, D.L., MacPhail, R.C., & Padilla, S. (2011). Rearing
538 conditions differentially affect the locomotor behavior of larval zebrafish, but not
539 their response to valproate-induced developmental neurotoxicity. *Neurotoxicology*
540 *and Teratology*, 33, 674-679.

541

542

Figure Legends

Figure 1. Schematic representation of experimental design. Grey boxes represent experimental units, text within black boxes indicates behavioural measurements and arrows represent progression of each treatment through time. Not to scale.

Figure 2. Behaviour of F_0 fish in the light and dark box test. (a) Time spent in the dark compartment and (b) activity, i.e. number of transitions between compartments for F_0 fish in the light and dark box test. Boxes represent the 25th and 75th percentiles and the line in between represents the median. Error bars are 95% confidence intervals. Sample sizes are given within the bars. Letters denote significant differences ($P < 0.05$), where bars sharing a letter are not significantly different.

Figure 3. Behaviour of F_0 fish in the novel tank diving test. (a) Latency to reach the upper compartment and (b) activity, i.e. number of transitions for F_0 fish in the novel tank diving test. Boxes represent the 25th and 75th percentiles and the line in between represents the median. Error bars are 95% confidence intervals. Sample sizes are given within the bars. Letters denote significant differences ($P < 0.05$), where bars sharing a letter are not significantly different.

Figure 4. Behaviour of F_0 and F_1 fish in the mirror image test. (a) Number of attacks/min in the mirror image test for F_0 fish, (b) number of attacks/min by offspring (F_1) and (c) time spent beside the mirror by offspring (F_1). Boxes represent the 25th and 75th percentiles and the line in between represents the median. Error bars are 95% confidence intervals. Sample sizes are given within the bars. Letters denote significant differences ($P < 0.05$), where bars sharing a letter are not significantly different.

Figure 1




Figure 2





Figure 3





Figure 4

